

Control Around Every Curve

Results

Patient 1

Diagnostic Ultrasound Images showed >25% decrease in thickness of the fat layer

Pre-treatment and 2-month post 2 treatments of subcutaneous fat showed a 0.64 cm reduction (2.44 cm to 1.80 cm) after truSculpt treatments. Courtesy of New Jersey Plastic Surgery Center, Barry



Abdomen

Patient 2

Pre-Treatment


4 Weeks Post 2 Treatments



Patient 3

Pre-Treatment

4 Weeks Post 2 Treatments



Patient 4



Treated side

Untreated Side

Patient 5



Pre-Treatment Post 2 treatments

Patient 6



Patient 7



Pre-Treatment


Post 2 treatments

Thigh


Before

4 Weeks Post 2 Treatments





Silhouette Overlay



IRB approved in-vivo clinical study performed at Cutera Research Center, Brisbane, CA USA - 2012

CUTERA © 2012 Cutera, Inc.

CE Mark cleared for body sculpting, body contouring, circumferential reduction, and fat reduction.

Cutera World Headquarters 3240 Bayshore Boulevard Brisbane, CA 94005 Telephone: (888) 4-CUTERA or (415) 657-5 Fax: (415) 330-3444

Sitemap | Privacy Policy